

$(\alpha')^4$ corrections to the $N = 2$ supersymmetric Born-Infeld action

A. De Giovanni ¹, A. Santambrogio ² and D. Zanon ¹

¹ Dipartimento di Fisica dell'Università di Milano
and INFN, Sezione di Milano, via Celoria 16, I-20133 Milano, Italy

² Instituut voor Theoretische Fysica, Katholieke Universiteit Leuven,
Celestijnenlaan 200-D, B-3001 Leuven, Belgium

ABSTRACT

We consider the $N = 2$ supersymmetric Born-Infeld action and compute one-loop divergences quantizing the theory in $N = 1$ superspace. We find that in the presence of non constant curvature the theory is not renormalizable. The structure of the $(\alpha')^4$ counterterm, proportional to derivatives of the curvature, is consistent with effective action calculations from superstring theory.

e-mail: antonio.degiovanni@mi.infn.it

e-mail: alberto.santambrogio@fys.kuleuven.ac.be

e-mail: daniela.zanon@mi.infn.it

July 1999

A nonlinear generalization of Maxwell theory is given by the Born-Infeld action [1]

$$S_{BI} = \frac{1}{\alpha^2} \int d^4x \left\{ 1 - \sqrt{-\det(\delta_{\mu\nu} + \alpha F_{\mu\nu})} \right\} \quad (1)$$

where we have defined $\alpha \equiv 2\pi\alpha'$ and have chosen a flat euclidean metric $g_{\mu\nu} = \delta_{\mu\nu}$. The action in (1) describes the self-interaction of electromagnetic fields of arbitrary degree in the curvature tensor $F_{\mu\nu}$. It has the remarkable property of reproducing the effective action from open bosonic strings in a background abelian gauge field with constant curvature [2, 3].

Given the fact that the Born-Infeld action contains terms of arbitrary order in the field-strength, a question that naturally arises is related to its renormalizability properties: is the theory finite, renormalizable or at least renormalizable for constant curvature? In the following we address these issues for the supersymmetric version of the theory, computing in superspace at the one-loop level.

The $N = 1$ supersymmetrization of the action in (1) has been obtained in ref. [4], while the extension to the $N = 2$ supersymmetric case has been presented in [5]. We focus on the $N = 2$ version of the theory: it can be obtained by a partial breaking of $N = 4$ supersymmetry and therefore it is consistent with the effective $D3$ -brane action [6] in the type IIB superstring theory.

We introduce the $N = 2$ superfield-strength $\mathcal{W}$ satisfying the constraints

$$\overline{D}_{\dot{\alpha}i}\mathcal{W} = 0 \quad D^4\mathcal{W} = \square\overline{\mathcal{W}} \quad (2)$$

with $i = \underline{1}, \underline{2}$. The $N = 2$ superspace invariant action can be written as [5]

$$S_{SBI} = \frac{1}{2} \int d^4x d^4\theta \mathcal{W}^2 + \frac{1}{8} \int d^4x d^4\theta d^4\overline{\theta} \mathcal{Y}(K, \overline{K}) \alpha^2 \mathcal{W}^2 \overline{\mathcal{W}}^2 \quad (3)$$

where $d^4\theta \equiv d^2\theta_{\underline{1}} d^2\theta_{\underline{2}}$. The function $\mathcal{Y}(K, \overline{K})$ defined in terms of $K = D^4\mathcal{W}^2$ is given explicitly by

$$\begin{aligned} \mathcal{Y}(K, \overline{K}) &= \frac{1 - \frac{\alpha^2}{4}(K + \overline{K}) - \sqrt{[1 - \frac{\alpha^2}{4}(K + \overline{K})]^2 - \frac{\alpha^4}{4}K\overline{K}}}{\frac{\alpha^4}{8}K\overline{K}} = \\ &= 1 + \frac{\alpha^2}{4}(K + \overline{K}) + \mathcal{O}(\alpha^4) \end{aligned} \quad (4)$$

Since we are interested in performing quantum calculations we rewrite the action in (3) in $N = 1$ superspace: there one can use the standard quantization procedure for

$N = 1$ superfields. To this end we decompose the $N = 2$ superfield-strength $\mathcal{W}$ in terms of its $N = 1$ superfield components

$$\begin{aligned}\mathcal{W} = & \Phi + \theta_{\underline{2}}^{\alpha} W_{\alpha} - \theta_{\underline{2}}^2 \bar{D}^2 \bar{\Phi} + \frac{i}{2} \theta_{\underline{2}}^{\alpha} \bar{\theta}_{\underline{2}}^{\dot{\alpha}} \partial_{\alpha\dot{\alpha}} \Phi + \\ & + \frac{i}{2} \theta_{\underline{2}}^2 \bar{\theta}_{\underline{2}}^{\dot{\alpha}} \partial_{\alpha\dot{\alpha}} W^{\alpha} + \frac{1}{4} \theta_{\underline{2}}^2 \bar{\theta}_{\underline{2}}^2 \square \Phi\end{aligned}\quad (5)$$

where $W_{\alpha} = i\bar{D}^2 D_{\alpha} V$ is the $N = 1$ chiral field-strength with V the unconstrained vector superfield, and Φ is the $N = 1$ matter superfield. In this manner the kinetic part of the action in (3) becomes (we define $\theta_{\perp\alpha} = \theta_{\alpha}$, $\bar{\theta}_{\perp\dot{\alpha}} = \bar{\theta}_{\dot{\alpha}}$)

$$\begin{aligned}S_{KIN} &= \frac{1}{2} \int d^4x d^4\theta \mathcal{W}^2 = \\ &= \int d^4x d^2\theta W^2 + \int d^4x d^2\theta d^2\bar{\theta} \Phi \bar{\Phi}\end{aligned}\quad (6)$$

The next contribution is given by the quartic interaction term $\mathcal{W}^2 \bar{\mathcal{W}}^2$ which can be easily decomposed in terms of $N = 1$ superfields

$$\begin{aligned}\mathcal{W}^2 \bar{\mathcal{W}}^2|_{\theta_{\underline{2}}^2 \bar{\theta}_{\underline{2}}^2} = & \frac{1}{4} \Phi^2 \square \bar{\Phi}^2 + \frac{1}{4} \bar{\Phi}^2 \square \Phi^2 + \frac{1}{4} i \partial^{\alpha\dot{\alpha}} (\Phi^2) i \partial_{\alpha\dot{\alpha}} (\bar{\Phi}^2) \\ & + 4\Phi \bar{D}^2 \bar{\Phi} \bar{\Phi} D^2 \Phi + 2\Phi \bar{D}^2 \bar{\Phi} \bar{W}^{\dot{\beta}} \bar{W}_{\dot{\beta}} + \\ & + 2W^{\beta} W_{\beta} \bar{\Phi} D^2 \Phi + W_{\beta} W^{\beta} \bar{W}_{\dot{\beta}} \bar{W}^{\dot{\beta}} + \\ & - 2i\Phi \bar{\Phi} W_{\beta} \partial^{\beta\dot{\beta}} \bar{W}_{\dot{\beta}} - 2i\Phi \partial^{\beta\dot{\beta}} \bar{\Phi} W_{\beta} \bar{W}_{\dot{\beta}} + \\ & + 2i\Phi \bar{\Phi} \partial^{\beta\dot{\beta}} W_{\beta} \bar{W}_{\dot{\beta}} + 2iW_{\beta} \partial^{\beta\dot{\beta}} \Phi \bar{\Phi} \bar{W}_{\dot{\beta}}\end{aligned}\quad (7)$$

The last two lines combine and up to total derivative terms the $\mathcal{O}(\alpha^2)$ contributions to the action are

$$\begin{aligned}S_{SBI}^{(\alpha^2)} &= \frac{\alpha^2}{8} \int d^4x d^4\theta d^4\bar{\theta} \mathcal{W}^2 \bar{\mathcal{W}}^2 = \\ &= \frac{\alpha^2}{2} \int d^4x d^2\theta d^2\bar{\theta} W^2 \bar{W}^2 - \frac{\alpha^2}{2} \int d^4x d^2\theta d^2\bar{\theta} \Phi \bar{W}^{\dot{\alpha}} i \partial_{\alpha\dot{\alpha}} (W^{\alpha} \Phi) \\ &+ \frac{\alpha^2}{2} \int d^4x d^2\theta d^2\bar{\theta} (\bar{W}^2 \Phi \bar{D}^2 \bar{\Phi} + W^2 \bar{\Phi} D^2 \Phi) + \\ &+ \frac{\alpha^2}{4} \int d^4x d^2\theta d^2\bar{\theta} \Phi i \partial^{\alpha\dot{\alpha}} \Phi \bar{\Phi} i \partial_{\alpha\dot{\alpha}} \bar{\Phi} + \\ &+ \frac{\alpha^2}{2} \int d^4x d^2\theta d^2\bar{\theta} \Phi \bar{D}^2 \bar{\Phi} \bar{\Phi} D^2 \Phi\end{aligned}\quad (8)$$

The other interactions from the expansion of $\mathcal{Y}(K, \overline{K})$ can be worked out straightforwardly, but the terms in (8) are sufficient for the determination of the one-loop counterterm. The gauge-fixing for the vector superfield action is standard (see for example [7]). We work in Feynman gauge and use the following propagators

$$\begin{aligned} \langle V(x, \theta) V(x', \theta') \rangle &= \frac{1}{\square} \delta^{(4)}(x - x') \delta^{(4)}(\theta - \theta') \\ \langle \Phi(x, \theta) \bar{\Phi}(x', \theta') \rangle &= -\frac{1}{\square} \delta^{(4)}(x - x') \delta^{(4)}(\theta - \theta') \end{aligned} \quad (9)$$

The vertices are obtained from (8) with extra $\overline{D}^2$ (D^2) factors acting on each chiral Φ (antichiral $\bar{\Phi}$) line leaving a vertex.

We are interested in the computation of $\mathcal{O}(\alpha^4)$ *on-shell* one-loop divergent contributions to the effective action and in particular we look for terms with external vector field-strengths. Once we have determined the structure of these counterterms, $N = 2$ supersymmetry will allow us to reconstruct the complete answer. Setting the external background on-shell means that we can freely use the equations of motion

$$D^\alpha W_\alpha = 0 \quad \overline{D}^{\dot{\alpha}} \overline{W}_{\dot{\alpha}} = 0 \quad (10)$$

which imply

$$D^2 W_\alpha = -D_\alpha D^\beta W_\beta = 0 \quad \overline{D}^2 \overline{W}_{\dot{\alpha}} = 0 \quad (11)$$

From the interaction terms in (8) we see that the relevant one-loop supergraphs are the ones with four external field-strengths and quantum propagators given by vector superfields and by matter chiral superfields. First we consider the diagrams with vector internal lines. Various different diagrams can be drawn: most of them are immediately discarded either because they trivially vanish after completion of the D -algebra, or are zero by momentum integration, or else they reduce to tadpole-type integrals which are set to zero in dimensional regularization scheme. The only non vanishing contributions correspond to the supergraphs shown in fig. 1 and 2 which we call G_1 and G_2 respectively in the following.

First we analyze the contribution to the effective action from G_1 . After D -algebra we obtain

$$\begin{aligned} \Gamma_{G_1}^{(4)}[W, \overline{W}] &= -\frac{\alpha^4}{32} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \bar{\theta} D^2 [W^\alpha(-p_1) W_\alpha(-p_2)] \\ &\quad \int \frac{d^{4-2\epsilon} k}{(2\pi)^4} \frac{(p+k)^{\beta\dot{\beta}} (-k)_{\beta\dot{\beta}}}{(p+k)^2 k^2} \overline{D}^2 [\overline{W}^{\dot{\alpha}}(-p_3) \overline{W}_{\dot{\alpha}}(-p_4)] \end{aligned} \quad (12)$$




Figure 1: Supegraph G_1 .




Figure 2: Supegraph G_2 .

where $p = p_1 + p_2 = -p_3 - p_4$ and momenta are labeled as shown in fig. 1.

Using dimensional regularization the momentum integral in (12) gives the following divergent contribution

$$\int \frac{d^{4-2\epsilon} k}{(2\pi)^4} \frac{(p+k)^{\alpha\dot{\alpha}} k_{\alpha\dot{\alpha}}}{(p+k)^2 k^2} \rightarrow -\frac{1}{\epsilon} \frac{p^2}{(4\pi)^2} \quad (13)$$

Therefore the divergence associated to the G_1 graph can be written as

$$\begin{aligned} \Gamma_{G_1}^{(4)} \text{div.} [W, \overline{W}] &= \frac{\alpha^4}{8} \left(-\frac{1}{\epsilon} \frac{1}{(4\pi)^2} \right) \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} \\ &\quad (p_1 + p_2)^2 W^2(1, 2) D^2 \overline{D}^2 \overline{W}^2(3, 4) \end{aligned} \quad (14)$$

In order to streamline the notation we write $W^2(1, 2) \equiv \frac{1}{2} W^\alpha(-p_1) W_\alpha(-p_2)$ and also $W_{\alpha,1} \equiv W_\alpha(-p_1)$ etc. Sometimes, whenever clear from the context, we simply drop the momentum suffixes.

It is useful to rewrite the result in (14) taking advantage of various on-shell identities. We list the ones that we repeatedly use

$$\begin{aligned}\overline{D}^2 \overline{W}^2 &= -\frac{1}{2} \left(\overline{D}^{\dot{\alpha}} \overline{W}^{\dot{\beta}} \right) \left(\overline{D}_{\dot{\alpha}} \overline{W}_{\dot{\beta}} \right) \\ D^2 \overline{D}^{\dot{\alpha}} \overline{W}^{\dot{\beta}} &= [D^2, \overline{D}^{\dot{\alpha}}] \overline{W}^{\dot{\beta}} = -p^{\alpha\dot{\alpha}} D_{\alpha} \overline{W}^{\dot{\beta}} = 0 \\ p^2 W_{\alpha}(p) &= -\overline{D}^2 D^2 W_{\alpha} = 0\end{aligned}\tag{15}$$

With the help of the above relations we can write

$$\begin{aligned}D^2 \overline{D}^2 \overline{W}^2(3, 4) &= -\frac{1}{2} \left[D^{\alpha} \overline{D}^{\dot{\alpha}} \overline{W}^{\dot{\beta}}(-p_3) \right] \left[D_{\alpha} \overline{D}_{\dot{\alpha}} \overline{W}_{\dot{\beta}}(-p_4) \right] = \\ &= -p_3^{\alpha\dot{\alpha}} p_4{}_{\alpha\dot{\alpha}} \overline{W}^2(3, 4) = -(p_3 + p_4)^2 \overline{W}^2(3, 4)\end{aligned}\tag{16}$$

The final result for the divergent contribution is

$$\begin{aligned}\Gamma_{G_1}^{(4)}{}_{div.} [W, \overline{W}] &= \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} \\ &\quad s^2 W^2(1, 2) \overline{W}^2(3, 4)\end{aligned}\tag{17}$$

We have introduced the Mandelstam variable $s = (p_1 + p_2)^2 = (p_3 + p_4)^2$. In short we write

$$\Gamma_{G_1}^{(4)}{}_{div.}(p_1, \dots, p_4) = \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} s^2\tag{18}$$

Now we turn to the corresponding calculation for the supergraph in fig. 2.

After completion of the D -algebra the G_2 diagram in fig. 2 gives rise to the following contribution to the effective action

$$\begin{aligned}\Gamma_{G_2}^{(4)} [W, \overline{W}] &= \frac{\alpha^4}{8} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} \overline{D}^{\dot{\gamma}} \left[D^{\gamma} W_{\alpha}(-p_1) \overline{W}_{\dot{\alpha}}(-p_3) \right] \\ &\quad \int \frac{d^{4-2\epsilon} k}{(2\pi)^4} \frac{(p+k)_{\gamma\dot{\gamma}} (p+k)^{\alpha\dot{\beta}} k^{\beta\dot{\alpha}}}{(p+k)^2 k^2} W_{\beta}(-p_2) \overline{W}_{\dot{\beta}}(-p_4)\end{aligned}\tag{19}$$

where now we have set $p = p_1 + p_3 = -p_2 - p_4$. The one-loop momentum integral leads to a divergence which in dimensional regularization is given by

$$\begin{aligned}
& \int \frac{d^{4-2\epsilon} k}{(2\pi)^4} \frac{(p+k)_{\gamma\dot{\gamma}} (p+k)^{\alpha\dot{\beta}} k^{\beta\dot{\alpha}}}{(p+k)^2 k^2} \\
& \rightarrow \frac{1}{12\epsilon} \frac{1}{(4\pi)^2} \left\{ -p_{\gamma\dot{\gamma}} p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} - p_{\gamma\dot{\gamma}} C^{\alpha\beta} C^{\dot{\beta}\dot{\alpha}} p^2 \right. \\
& \quad \left. + p^{\beta\dot{\alpha}} p^2 \delta_{\gamma}^{\alpha} \delta_{\dot{\gamma}}^{\dot{\beta}} - p^{\alpha\dot{\beta}} p^2 \delta_{\gamma}^{\beta} \delta_{\dot{\gamma}}^{\dot{\alpha}} \right\}
\end{aligned} \tag{20}$$

so that we obtain

$$\begin{aligned}
\Gamma_{G_2}^{(4)} \text{div} [W, \overline{W}] &= \frac{\alpha^4}{96} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} \\
& \left\{ -p_{\gamma\dot{\gamma}} p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} [\overline{D}^{\dot{\gamma}} D^{\gamma} (W_{\alpha,1} \overline{W}_{\dot{\alpha},3})] W_{\beta,2} \overline{W}_{\dot{\beta},4} \right. \\
& + p_{\gamma\dot{\gamma}} p^2 [\overline{D}^{\dot{\gamma}} D^{\gamma} (W_1^{\alpha} \overline{W}_3^{\dot{\alpha}})] W_{\alpha,2} \overline{W}_{\dot{\alpha},4} \\
& + p^{\beta\dot{\alpha}} p^2 [\overline{D}^{\dot{\beta}} D^{\alpha} (W_{\alpha,1} \overline{W}_{\dot{\alpha},3})] W_{\beta,2} \overline{W}_{\dot{\beta},4} \\
& \left. - p^{\alpha\dot{\beta}} p^2 [\overline{D}^{\dot{\alpha}} D^{\beta} (W_{\alpha,1} \overline{W}_{\dot{\alpha},3})] W_{\beta,2} \overline{W}_{\dot{\beta},4} \right\}
\end{aligned} \tag{21}$$

Now we impose the on-shell equations (10,11,15) on the external background. In addition we use the following relations

$$p_{\gamma\dot{\gamma}} (D^{\gamma} W_{\alpha,1}) (\overline{D}^{\dot{\gamma}} \overline{W}_{\dot{\alpha},3}) = (p_1 + p_3)_{\gamma\dot{\gamma}} (D^{\gamma} W_{\alpha,1}) (\overline{D}^{\dot{\gamma}} \overline{W}_{\dot{\alpha},3}) = 0 \tag{22}$$

Also for two spinors $\psi_{\alpha}, \chi_{\alpha}$ we can write

$$\psi_{\alpha} \chi_{\beta} = \frac{1}{2} \psi_{(\alpha} \chi_{\beta)} + \frac{1}{2} C_{\beta\alpha} \psi^{\gamma} \chi_{\gamma} \tag{23}$$

In this way we have

$$\begin{aligned}
& p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} W_{\alpha,1} \overline{W}_{\dot{\alpha},3} W_{\beta,2} \overline{W}_{\dot{\beta},4} = \\
& = -p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} \left[\frac{1}{2} (W_{\alpha,1} W_{\beta,2} + W_{\beta,1} W_{\alpha,2}) + \frac{1}{2} C_{\beta\alpha} W_1^{\gamma} W_{\gamma,2} \right] \\
& \quad \left[\frac{1}{2} (\overline{W}_{\dot{\alpha},3} \overline{W}_{\dot{\beta},4} + \overline{W}_{\dot{\beta},3} \overline{W}_{\dot{\alpha},4}) + \frac{1}{2} C_{\dot{\beta}\dot{\alpha}} \overline{W}_3^{\dot{\gamma}} \overline{W}_{\dot{\gamma},4} \right] \\
& = -\frac{1}{2} p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} W_{\alpha,1} W_{\beta,2} \overline{W}_{\dot{\alpha},3} \overline{W}_{\dot{\beta},4} - C_{\beta\alpha} C_{\dot{\beta}\dot{\alpha}} W^2(1,2) \overline{W}^2(3,4) p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}}
\end{aligned} \tag{24}$$

since on-shell

$$\begin{aligned} p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} W_{\alpha,1} W_{\beta,2} \overline{W}_{\dot{\beta},3} \overline{W}_{\dot{\alpha},4} &= 0 \\ p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} (W_{\alpha,1} W_{\beta,2} + W_{\beta,1} W_{\alpha,2}) C_{\dot{\beta}\dot{\alpha}} \overline{W}_3^{\dot{\gamma}} \overline{W}_{\dot{\gamma},4} &= 0 \end{aligned} \quad (25)$$

Thus (24) leads to the on-shell identity

$$\begin{aligned} p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} W_{\alpha,1} \overline{W}_{\dot{\alpha},3} W_{\beta,2} \overline{W}_{\dot{\beta},4} &= \\ = -2 C_{\beta\alpha} C_{\dot{\beta}\dot{\alpha}} W^2(1,2) \overline{W}^2(3,4) p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} &= \\ = 4 p^2 W^2(1,2) \overline{W}^2(3,4) \end{aligned} \quad (26)$$

Finally we obtain

$$\begin{aligned} \Gamma_{G_2 \text{ div}}^{(4)} [W, \overline{W}] &= -\frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \frac{1}{12} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} \\ &\quad \left\{ p_{\gamma\dot{\gamma}} p^{\alpha\dot{\beta}} p^{\beta\dot{\alpha}} (-p_1^{\gamma\dot{\gamma}}) - p_{\gamma\dot{\gamma}} C^{\alpha\beta} C^{\dot{\alpha}\dot{\beta}} p^2 (-p_1^{\gamma\dot{\gamma}}) + p^{\alpha\dot{\beta}} p^2 (-p_1^{\beta\dot{\alpha}}) \right\} \\ &\quad W_{\alpha}(-p_1) \overline{W}_{\dot{\alpha}}(-p_3) W_{\beta}(-p_2) \overline{W}_{\dot{\beta}}(-p_4) = \\ &= \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} (p^2)^2 W^2(1,2) \overline{W}^2(3,4) \end{aligned} \quad (27)$$

In short we write

$$\Gamma_{G_2 \text{ div}}^{(4)}(p_1, \dots, p_4) = \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{t^2}{(4\pi)^2} \quad (28)$$

where $t = (p_1 + p_3)^2 = (p_2 + p_4)^2$.

Now we study the contributions from the diagrams with chiral internal lines. Setting the external vector background on-shell, the relevant vertex from the action in (8) is $\frac{\alpha^2}{2} \int d^4 x d^2 \theta d^2 \overline{\theta} W_{\alpha} \overline{W}_{\dot{\alpha}} \Phi i \partial^{\alpha\dot{\alpha}} \Phi$. Thus we obtain a third graph that we call G_3 . It is straightforward to realize that the D-algebra is the same as the one for the G_2 diagram shown in fig. 2, with $\dot{\alpha} \leftrightarrow \dot{\beta}$ in the loop. Repeating exactly the same steps as for G_2 , we obtain

$$\begin{aligned} \Gamma_{G_3 \text{ div}}^{(4)} [W, \overline{W}] &= \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \frac{1}{12} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} \\ &\quad p_{\gamma\dot{\gamma}} C^{\beta\alpha} C^{\dot{\beta}\dot{\alpha}} p^2 (-p_1^{\gamma\dot{\gamma}}) W_{\alpha}(-p_1) \overline{W}_{\dot{\alpha}}(-p_3) W_{\beta}(-p_2) \overline{W}_{\dot{\beta}}(-p_4) = \\ &= \frac{\alpha^4}{24} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \overline{\theta} (p^2)^2 W^2(1,2) \overline{W}^2(3,4) \end{aligned} \quad (29)$$

The total one-loop divergence for $W^2\overline{W}^2$ is given by the sum of (17), (27) and (29)

$$\Gamma_{div}^{(4)}[W, \overline{W}] = \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \int d^4 p_1 \dots d^4 p_4 \delta(\Sigma p_i) d^2 \theta d^2 \bar{\theta} (s^2 + \frac{4}{3}t^2) W^2(1, 2) \overline{W}^2(3, 4) \quad (30)$$

The $N = 2$ supersymmetrization of the result is immediate since we have shown that the term $W^2\overline{W}^2$ appears in the full $N = 2$ superspace integral of $\mathcal{W}^2\overline{\mathcal{W}}^2$ (see eq. (7)). In fact, using the equations of motion for the vectors and for the matter fields ($D^2\Phi = 0$, $\overline{D}^2\overline{\Phi} = 0$), the only terms that survive on-shell are

$$\frac{\alpha^2}{2} \int d^4 x d^2 \theta d^2 \bar{\theta} \left(W^2\overline{W}^2 + W_\alpha \overline{W}_{\dot{\alpha}} \overline{\Phi} i \partial^{\alpha\dot{\alpha}} \Phi - \frac{1}{4} \overline{\Phi}^2 i \partial^{\alpha\dot{\alpha}} \Phi i \partial_{\alpha\dot{\alpha}} \Phi \right) \quad (31)$$

The $\mathcal{O}(\alpha^4)$ counterterm given in (30) can be rewritten in configuration space

$$\Gamma_{div}^{(4)} = \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \int d^4 x d^2 \theta d^2 \bar{\theta} (\partial^\mu W^\alpha \partial_\mu W_\alpha \partial^\nu \overline{W}^{\dot{\alpha}} \partial_\nu \overline{W}_{\dot{\alpha}} + \frac{4}{3} \partial^\mu W^\alpha \partial^\nu W_\alpha \partial_\mu \overline{W}^{\dot{\alpha}} \partial_\nu \overline{W}_{\dot{\alpha}}) \quad (32)$$

In order to make contact with the standard bosonic Born-Infeld action, we express the result in component form. The θ -expansion of the superfield-strength is

$$W_\alpha(x, \theta, \bar{\theta}) = \lambda_\alpha + \theta^\beta f_{\alpha\beta} - i\theta^\beta C_{\beta\alpha} D' + i\theta^2 \partial_{\alpha\dot{\alpha}} \overline{\lambda}^{\dot{\alpha}} + \frac{i}{2} \theta^\beta \bar{\theta}^{\dot{\beta}} \partial_{\beta\dot{\beta}} \lambda_\alpha + \frac{1}{4} \theta^2 \bar{\theta}^2 \square \lambda_\alpha - \frac{i}{2} \theta^2 \bar{\theta}^{\dot{\beta}} \partial_{\dot{\beta}}^\beta (f_{\alpha\beta} - i C_{\beta\alpha} D') \quad (33)$$

$$\overline{W}_{\dot{\alpha}}(x, \theta, \bar{\theta}) = \overline{\lambda}_{\dot{\alpha}} + \bar{\theta}^{\dot{\beta}} (\overline{f}_{\dot{\alpha}\dot{\beta}} + i C_{\dot{\beta}\dot{\alpha}} D') + i \bar{\theta}^2 \partial_{\alpha\dot{\alpha}} \lambda^\alpha + \frac{i}{2} \bar{\theta}^{\dot{\beta}} \theta^\beta \partial_{\beta\dot{\beta}} \overline{\lambda}_{\dot{\alpha}} + \frac{1}{4} \theta^2 \bar{\theta}^2 \square \overline{\lambda}_{\dot{\alpha}} - \frac{i}{2} \bar{\theta}^2 \theta^\gamma \partial_\gamma^{\dot{\beta}} (\overline{f}_{\dot{\alpha}\dot{\beta}} + i C_{\dot{\beta}\dot{\alpha}} D') \quad (34)$$

where $f_{\alpha\beta}$ (similarly $\overline{f}_{\dot{\alpha}\dot{\beta}}$) is related to $F_{\mu\nu}$ by

$$\begin{aligned} f_{\alpha\beta} &\equiv \frac{1}{2} (\sigma^{\mu\nu})_{\alpha\beta} F_{\mu\nu} \\ (\sigma_{\mu\nu})_\alpha^\beta &= -\frac{1}{4} (\sigma_\mu \tilde{\sigma}_\nu - \sigma_\nu \tilde{\sigma}_\mu)_\alpha^\beta \\ \sigma_\mu &= (1, \vec{\sigma}) \quad \tilde{\sigma}_\mu = (1, -\vec{\sigma}) \end{aligned} \quad (35)$$

Thus we can easily obtain the $\theta^2 \bar{\theta}^2$ coefficient of the component expansion of $W^2 \bar{W}^2$

$$\begin{aligned}
& W^\alpha W_\alpha \bar{W}^{\dot{\alpha}} \bar{W}_{\dot{\alpha}}|_{\theta^2 \bar{\theta}^2} = \\
& = \frac{1}{4} \lambda^\alpha \lambda_\alpha \square (\bar{\lambda}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}}) + \frac{1}{4} \square (\lambda^\alpha \lambda_\alpha) \bar{\lambda}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}} + \frac{1}{4} i \partial^{\beta \dot{\beta}} (\lambda^\alpha \lambda_\alpha) i \partial_{\beta \dot{\beta}} (\bar{\lambda}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}}) \\
& \quad + 2i \partial^{\beta \dot{\beta}} [\lambda^\alpha (f_{\alpha\beta} - i C_{\beta\alpha} D')] \bar{\lambda}^{\dot{\alpha}} (\bar{f}_{\dot{\alpha}\beta} + i C_{\beta\dot{\alpha}} D') + \\
& \quad + 2i \partial^{\beta \dot{\beta}} [\bar{\lambda}^{\dot{\alpha}} (\bar{f}_{\dot{\alpha}\beta} + i C_{\beta\dot{\alpha}} D')] \lambda^\alpha (f_{\alpha\beta} - i C_{\beta\alpha} D') - \\
& \quad - 4 \lambda^\alpha \partial_{\alpha\dot{\alpha}} \bar{\lambda}^{\dot{\alpha}} \bar{\lambda}^{\dot{\beta}} \partial_{\beta\dot{\beta}} \lambda^\beta + \\
& \quad + 2i \lambda^\gamma \partial_{\gamma\dot{\gamma}} \bar{\lambda}^{\dot{\gamma}} (\bar{f}^{\dot{\alpha}\dot{\beta}} \bar{f}_{\dot{\alpha}\dot{\beta}} + 2i \bar{f}^{\dot{\alpha}}_{\dot{\alpha}} D' - 2(D')^2) + \\
& \quad + 2i (f^{\alpha\beta} f_{\alpha\beta} - 2i f^\alpha_\alpha D' - 2(D')^2) \bar{\lambda}^{\dot{\gamma}} \partial_{\gamma\dot{\gamma}} \lambda^\gamma + \\
& \quad + (f^{\alpha\beta} f_{\alpha\beta} - 2i f^\alpha_\alpha D' - 2(D')^2) (\bar{f}^{\dot{\alpha}\dot{\beta}} \bar{f}_{\dot{\alpha}\dot{\beta}} + 2i \bar{f}^{\dot{\alpha}}_{\dot{\alpha}} D' - 2(D')^2) \quad (36)
\end{aligned}$$

This expression can be simplified using the equations of motion for the auxiliary field D' . One can easily show that to all orders in α an on-shell solution is given by $D' = 0$. In this way we obtain

$$\begin{aligned}
& W^\alpha W_\alpha \bar{W}^{\dot{\alpha}} \bar{W}_{\dot{\alpha}}|_{\theta^2 \bar{\theta}^2, D'=0} = \\
& = \frac{1}{4} \lambda^\alpha \lambda_\alpha \square (\bar{\lambda}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}}) + \frac{1}{4} \square (\lambda^\alpha \lambda_\alpha) \bar{\lambda}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}} + \frac{1}{4} i \partial^{\beta \dot{\beta}} (\lambda^\alpha \lambda_\alpha) i \partial_{\beta \dot{\beta}} (\bar{\lambda}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}}) \\
& \quad + 2i \partial^{\beta \dot{\beta}} (\lambda^\alpha f_{\alpha\beta}) \bar{\lambda}^{\dot{\alpha}} \bar{f}_{\dot{\alpha}\beta} + 2i \partial^{\beta \dot{\beta}} (\bar{\lambda}^{\dot{\alpha}} \bar{f}_{\dot{\alpha}\beta}) \lambda^\alpha f_{\alpha\beta} - \\
& \quad - 4 \lambda^\alpha \partial_{\alpha\dot{\alpha}} \bar{\lambda}^{\dot{\alpha}} \bar{\lambda}^{\dot{\beta}} \partial_{\beta\dot{\beta}} \lambda^\beta + \\
& \quad + 2i \lambda^\gamma \partial_{\gamma\dot{\gamma}} \bar{\lambda}^{\dot{\gamma}} \bar{f}^{\dot{\alpha}\dot{\beta}} \bar{f}_{\dot{\alpha}\dot{\beta}} + 2i f^{\alpha\beta} f_{\alpha\beta} \bar{\lambda}^{\dot{\gamma}} \partial_{\gamma\dot{\gamma}} \lambda^\gamma + \\
& \quad + f^{\alpha\beta} f_{\alpha\beta} \bar{f}^{\dot{\alpha}\dot{\beta}} \bar{f}_{\dot{\alpha}\dot{\beta}} \quad (37)
\end{aligned}$$

Going back to (32) we obtain

$$\begin{aligned}
& \Gamma_{div}^{(4)}[W, \bar{W}] = \Gamma_{div}^{(4)}[F^+, F^-] + \dots = \\
& = \frac{\alpha^4}{8} \frac{1}{\epsilon} \frac{1}{(4\pi)^2} \int d^4 x \left[\partial^\mu (F^+)^{\rho\sigma} \partial_\mu (F^+)_{\rho\sigma} \partial^\nu (F^-)^{\rho'\sigma'} \partial_\nu (F^-)_{\rho'\sigma'} + \right. \\
& \quad \left. + \frac{4}{3} \partial^\mu (F^+)^{\rho\sigma} \partial^\nu (F^+)_{\rho\sigma} \partial_\mu (F^-)^{\rho'\sigma'} \partial_\nu (F^-)_{\rho'\sigma'} \right] + \dots \quad (38)
\end{aligned}$$

where we have introduced the definitions

$$\tilde{F}_{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu}^{\rho\sigma} F_{\rho\sigma} \quad F^{(\pm)} = \frac{1}{2} (F \pm i\tilde{F}) \quad (39)$$

and used the relation

$$f^{\alpha\beta} f_{\alpha\beta} \overline{f}^{\dot{\alpha}\dot{\beta}} \overline{f}_{\dot{\alpha}\dot{\beta}} = (F^+)^2 (F^-)^2 \quad (40)$$

In (40) we have dropped total derivative terms that anyway would not contribute upon d^4x integration.

We can rewrite the $\Gamma^{(4)}[F]$ contribution in momentum space: using

$$(F^+)^2 (F^-)^2 = F^4 - \frac{1}{4} (F^2)^2 \quad (41)$$

with

$$\begin{aligned} F\tilde{F} &= F^{\mu\nu} \tilde{F}_{\mu\nu} & F^2 &= F^{\mu\nu} F_{\mu\nu} \\ F^4 &= F^{\mu\nu} F_{\nu\lambda} F^{\lambda\rho} F_{\rho\mu} = \frac{1}{4} (F\tilde{F})^2 + \frac{1}{2} (F^2)^2 \end{aligned} \quad (42)$$

we obtain

$$\Gamma_{div}^{(4)}[F] = \frac{\alpha^4}{32\epsilon} \frac{1}{(4\pi)^2} \int d^4p_1 \dots d^4p_4 \delta(\Sigma p_i) (s^2 + \frac{4}{3}t^2) \left[F^4 - \frac{1}{4} (F^2)^2 \right] \quad (43)$$

Finally, since the factor $F^4 - \frac{1}{4} (F^2)^2$ is completely symmetric in the s, t, u variables, the result in (43) can be rewritten as

$$\Gamma_{div}^{(4)}[F] = \frac{7}{6} \frac{\alpha^4}{48\epsilon} \frac{1}{(4\pi)^2} \int d^4p_1 \dots d^4p_4 \delta(\Sigma p_i) (s^2 + t^2 + u^2) \left[F^4 - \frac{1}{4} (F^2)^2 \right] \quad (44)$$

It is interesting to note that terms like the one in eq. (44) have been obtained in the past from the study of scattering amplitudes of vector fields in type IIB string theory on the $D3$ -brane [8] and in type I open string theory [9, 10, 11]. Indeed the result obtained from string theory is of the form (cfr. [8])

$$\frac{\Gamma(s)\Gamma(t)}{\Gamma(1+s+t)} K(\epsilon^{(i)}, p^{(j)}) \quad (45)$$

where $\epsilon^{(i)}$ are the four polarization vectors and $p^{(j)}$ are the four external momenta. K is a kinematic factor corresponding to the term $\left[F^4 - \frac{1}{4}(F^2)^2\right]$. As mentioned above this factor is completely symmetric in the external momenta. Expanding the Γ -functions in eq. (45) and symmetrizing with respect to the momenta we obtain a term proportional to

$$(s^2 + t^2 + u^2) \left[F^4 - \frac{1}{4}(F^2)^2 \right], \quad (46)$$

i.e. of the same form as in (44).

The result in (44) gives the $\mathcal{O}(\alpha^4)$ counterterm of the bosonic Born-Infeld action. On dimensional grounds it is easy to determine the general structure of the allowed counterterms at any order. Taking into account the mass dimensions of the various factors

$$d^4x \sim m^{-4} \quad \alpha \sim m^{-2} \quad \partial_\mu \sim m \quad F_{\mu\nu} \sim m^2 \quad (47)$$

and keeping in mind that divergences correspond to local expressions, we have that the structure of the counterterms at L -loop order can be written symbolically as

$$\frac{1}{\alpha^2} \int d^4x (\alpha^2 \partial^4)^L (\alpha F)^{2n} \quad n > 1 \quad (48)$$

It is clear that in the presence of constant curvature F no divergent correction arises. On the contrary, whenever ∂F is not negligible divergences might appear at any loops and to all orders in the fields. The number of derivatives is fixed ∂^{4L} at L loops, while any number of fields can be produced since αF is of mass dimensions zero.

In the same way one can repeat this analysis for the supersymmetric theory. It would be quite interesting to perform higher order computations explicitly and determine the counterterms exactly.

In a recent paper [12] similar issues have been addressed and one-loop corrections have been computed for the $D3$ brane action.

Acknowledgments

We thank Renata Kallosh for having raised our interest on the subject.

This work was supported by the European Commission TMR program ERBFMRX-CT96-0045, in which A. DG. and D. Z. are associated to the University of Torino.

References

- [1] M. Born, L. Infeld, Proc. R. Soc. **A144** (1934) 425
- [2] E.S. Fradkin, A.A. Tseytlin, Phys. Lett. **B163** (1985) 123
- [3] A. Abouelsaood, C.G.Callan, C.R. Nappi, S.A. Yost, Nucl. Phys. **B280** (1987) 599
- [4] S. Cecotti, S. Ferrara, Phys. Lett. **187B** (1987) 335
- [5] S.V. Ketov, Mod. Phys. Lett. **A14** (1999) 501, hep-th/9809121; "Born-Infeld-Goldstone superfield action for gauge-fixed D-5- and D-3-branes in 6d", hep-th/9812051
- [6] M. Aganin, C. Popescu, J.H. Schwarz, Nucl. Phys. **B495** (1997) 99
- [7] S.J. Gates, M.T. Grisaru, M. Rocek, W. Siegel, "Superspace or one thousand and one lessons in supersymmetry", Benjamin Cummings, Reading MA, 1983
- [8] A. Hashimoto and I.R. Klebanov, Phys. Lett. **B381** (1996) 437, hep-th/9604065; Nucl. Phys. Proc. Suppl.**55B** (1997) 118, hep-th/9611214
- [9] J.H. Schwarz, Phys. Rep. **89** (1982) 223
- [10] D.J. Gross and E. Witten, Nucl. Phys. **B277** (1986) 1
- [11] O.D. Andreev and A.A. Tseytlin, Nucl. Phys. **B311** (1988) 205
- [12] M. Shmakova, "One-loop corrections to the $D3$ brane action", hep-th/9906239.